



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

arc-light directed upon it at the same time, the lighter portions of the powder were repelled, producing an appearance closely resembling the tail of a comet. The effect was comparable with what was to be expected from the energy observed. The authors considered it possible that the phenomena might be produced in part by other causes. In any event, the experiment reproduced with great accuracy the appearances observed in the tails of comets.

This work of NICHOLS and HULL points the way to other investigations which should throw much light on the vexed question of the constitution of the comets. C. D. P.

NOTES ON THE THREE COMETS NOW VISIBLE.

Comet a 1903: The telegram announcing the discovery of a new comet by M. GIACOBINI at Nice was received at the Lick Observatory on January 20th, but continuous stormy weather prevented observations here until the evening of January 29th. As seen with the 12-inch telescope on that date, the comet appeared to be about as bright as a ninth-magnitude star. It was small, nearly round, and with only a feeble condensation. During the month of February the comet grew steadily brighter and larger, and the condensation in the coma became more sharply defined, appearing at the time of the last observation as a bright disk about 2" or 3" in diameter. On February 15th, a faint straight tail was visible to a distance of about 5' in the north-following quadrant. This has grown brighter and longer, being traced to a distance of more than one degree in position-angle 47° on February 25th. When last seen the comet as a whole was as bright as a 6.5-magnitude star.

Comet d 1902: Observations secured here during February show that this comet is following very closely the orbit given in No. 88 of these *Publications* (p. 27). The residuals at the date of my last observations (February 28th) were only $-0^s.6$ and $+9''$. The comet's appearance has changed very little, though it is perhaps fainter than at first. It is still measurable with the 12-inch telescope, and can probably be followed with large telescopes for a month or two more. With one exception its perihelion distance is greater than that of any other comet so far found.

Comet b 1902: PERRINE'S comet was last observed here before its perihelion passage on November 17, 1902. After passing the Sun the comet was visible only to observers in the southern hemisphere until late in January, 1903. On the 29th of January, I found the comet close to the place predicted by STRÖMGREN'S ephemeris, the observed data being:—

$$\begin{aligned}\text{Jan. 29, } 12^{\text{h}} 57^{\text{m}} 38^{\text{s}} \text{ Mt. Ham. M. T. } \alpha &= 9^{\text{h}} 5^{\text{m}} 53^{\text{s}}.99 \\ \delta &= -41^{\circ} 0' 55''.7\end{aligned}$$

Since then the comet has moved rapidly north and west, passing close to *Sirius* in the latter part of February. On February 28th it was still visible in the 3-inch finder, and the 36-inch showed a faint nucleus.

Stormy weather has prevented observations of any of these comets since February 28th.

March 16, 1903.

R. G. AITKEN.

ASTRONOMICAL TELEGRAMS.

(Translations.)

CAMBRIDGE, MASS., Jan. 20, 1903.

To Lick Observatory: (Received 8:30 A. M.)

A comet was discovered by GIACOBINI at Nice on January 15th. It was observed at Nice on January 19.2498 G. M. T., in R. A. $22^{\text{h}} 57^{\text{m}} 48^{\text{s}}.0$; Decl. $+2^{\circ} 17' 36''$. The daily motion is east $17'$, north $12'$. (Signed) E. C. PICKERING.

BOSTON, MASS., Jan. 22, 1903.

To Lick Observatory: (Received 8:30 A. M.)

Comet *a 1903* (GIACOBINI) was observed by DINWIDDIE at Washington, on January 21.4915 G. M. T., in R. A. $23^{\text{h}} 0^{\text{m}} 6^{\text{s}}.5$; Decl. $+2^{\circ} 47' 46''$. (Signed) JOHN RITCHIE, JR.